



A Water Mass Classification Approach to Tracking Variability in the East Australian Current

Lachlan R. Phillips^{1*}, Gemma Carroll^{2,3}, Ian Jonsen¹, Robert Harcourt¹ and Moninya Roughan⁴

¹ Department of Biological Sciences, Macquarie University, Sydney, NSW, Australia, ² Institute of Marine Sciences, University of California, Santa Cruz, Santa Cruz, CA, United States, ³ Environmental Research Division, NOAA Southwest Fisheries Science Center, Monterey, California, CA, United States, ⁴ School of Mathematics and Statistics, UNSW Sydney, Sydney, NSW, Australia

OPEN ACCESS

Edited by:

Katrin Schroeder,
Italian National Research Council
(CNR), Italy

Reviewed by:

Antonio Olita,
Italian National Research Council
(CNR), Italy
Manuel Bensi,
Istituto Nazionale di Oceanografia e di
Geofisica Sperimentale (OGS), Italy

*Correspondence:

Lachlan R. Phillips
lachlan.phillips@hdr.mq.edu.au

Specialty section:

This article was submitted to
Physical Oceanography,
a section of the journal
Frontiers in Marine Science

Received: 27 February 2020

Accepted: 29 April 2020

Published: 03 June 2020

Citation:

Phillips LR, Carroll G, Jonsen I,
Harcourt R and Roughan M (2020) A
Water Mass Classification Approach
to Tracking Variability in the East
Australian Current.
Front. Mar. Sci. 7:365.
doi: 10.3389/fmars.2020.00365

The East Australian Current (EAC) is a southward flowing western boundary current that transports relatively warm and nutrient-depleted subtropical water along Australia's east coast. The EAC is a highly variable system that is formed by temporally-varying mixtures of water in the Coral Sea that do not form a linear density gradient or conform to a set range of temperature and salinity values. It can therefore be difficult to track EAC dynamics across both space and time using traditional analytical approaches. In order to more accurately quantify variability and trends in penetration of the EAC we develop a novel machine-learning classification approach to quantify variability in coastal EAC dynamics along a latitudinal gradient within the EAC extension zone in southeastern Australia. Applying our method to data from a 22-year free running regional hydrodynamic model revealed significant decadal-scale changes to EAC dynamics in the region. The annual period (generally in the austral summer) when the EAC is the dominant water mass in the region increased by approximately 2 months over the model time series. The encroachment of the EAC's traditional period of summer dominance into winter may have significant ecological implications through the acceleration of poleward range extensions by vagrant tropical species, facilitation of community phase shifts from temperate to tropical assemblages, and a phenological shift in the timing of major phytoplankton blooms. These results highlight the need to further understand the rapid changes occurring within western boundary current systems, and illustrates how classification approaches may assist in uncovering patterns in these highly variable systems.

Keywords: western boundary current, east Australian current, classification, water mass, EAC extension, intensification, current dynamics

1. INTRODUCTION

Western boundary currents are strong and persistent features of ocean circulation that form along the western margins of the world's major ocean basins (Hogg and Johns, 1995; Imawaki et al., 2013). Primarily generated by wind-driven anticyclonic subtropical gyres, these currents play a key role in meridional heat transport and climate regulation through the transportation of warm subtropical waters from low to high latitudes (Yu and Weller, 2007). Due to baroclinic instabilities, western boundary currents have characteristically elevated levels of eddy kinetic energy (Archer et al., 2018). As a consequence, western boundary currents are typically highly variable and dynamic, making it difficult to track these systems through time.

Because the water mass does not typically form a linear density gradient or conform to a set range of temperature and salinity values, it can be difficult to determine when a given region is experiencing flow originating predominately from the EAC or from other regions.

Machine learning is a tool that can be leveraged to classify dynamic and complex features in oceanic systems (Jones et al., 2014; Gangopadhyay et al., 2015). Machine learning algorithms are designed to learn from one dataset (training set) to make predictions about a new, independent dataset (testing set) and are well suited to analysing big datasets common in oceanographic studies. Multiple machine learning techniques are available to approach classification problems (Weiss and Kulikowski, 1991; Kotsiantis, 2007) including logistic regression which is a particularly well suited tool to model the outcomes of categorical dependent variables (Hosmer et al., 2013) such as the presence/absence of ocean currents.

This study describes variability in EAC dynamics along a latitudinal gradient within the EAC extension, downstream of its typical separation zone, along southeastern Australia. This is achieved through the development of a novel logistic regression classification algorithm that can identify water as being either subtropical origin (STO) or Tasman Sea origin (TSO). Classification is based on the surface temperature and salinity signatures of the two primary water masses known to influence our study region as proxies for EAC presence/absence. We then apply the classification algorithm to the output of a 22-year regional hydrodynamic model (Kerry and Roughan, 2020a) to examine variability and trends in EAC influence through time. We aim to provide a more detailed description of seasonal and inter-annual EAC dynamics, as well as describing spatial and temporal patterns of intensification in one of the fastest warming regions in the Southern Hemisphere. Our approach offers a framework to analyse variability in water mass influence that can be applied to other ocean systems.

2. METHODS

2.1. Study Region

To assess latitudinal variability in EAC dynamics along Australia's east coast, we analyzed variability in EAC presence for three zones (100 km squares) centered on coastal marine park ecosystems: (i) Jervis Bay Marine Park (35.114°S, 150.95°E); (ii) Batemans Marine Park (36.227°S, 150.23°E); and (iii) Cape Howe Marine Park (37.587°S, 150.29°E) (Figure 1). These locations were selected as they each encompass coastal ecosystems of conservation concern that lie in the path of the EAC extension. Each study site was restricted to the area between the coastal boundary and the 4,000 m depth contour.

2.2. Data

The classification algorithm utilized surface temperature and salinity data from two sources: (i) the “CSIRO Atlas of Regional Seas” (CARS) (Ridgway et al., 2015) as the training set and; (ii) a custom configuration of the “Regional Ocean Modeling System”

(ROMS 3.4) for circulation along the southeast coast of Australia (Kerry and Roughan, 2020b) as the testing set.

The CARS dataset is an atlas of ocean water properties derived from a quality-controlled archive of ocean measurements taken from research vessel instrument profiles and Argo floats from 1985 to 2009. These measurements are gridded at a 0.5 degree grid spatial resolution and averaged through time to provide a daily climatology of ocean conditions for each grid cell (Ridgway et al., 2002).



ROMS is a free-surface, hydrostatic, primitive equation ocean model solved on a curvilinear grid with a terrain following a vertical coordinate system (Shchepetkin and McWilliams, 2003, 2005). Time-stamped surface temperature and salinity data at a horizontal spatial resolution of ~2 km were extracted from a configuration of ROMS that produces a high resolution free running simulation of the EAC region (Kerry and Roughan, 2020a). Data were extracted for the period from 1994 to mid 2016. The BlueLink Reanalysis (BRAN3p5) (Oke et al., 2008) is used for the EAC-ROMS model initial conditions and boundary forcing over the 22-year period. EAC-ROMS has been validated against satellite and *in situ* observations to confirm that it resolves the mean dynamic features of the EAC including surface and subsurface variability, seasonality, and to ensure that it contains no net temperature drift (Kerry and Roughan, 2020a). For a full description of the EAC-ROMS specifications, parameters, and validation (see Kerry and Roughan, 2020a,b).

2.3. Algorithm Construction and Analysis

To quantify variability in EAC dynamics for each of the study sites, a supervised machine-learning classification algorithm was developed to assess the probability of surface water from the ROMS model output as being of either subtropical origin (STO) or Tasman Sea origin (TSO) based on the surface temperature and salinity signatures as a proxy for EAC-derived/non-EAC derived water. To do this, we undertook the following steps (Figure 2):

(1) The classification algorithm was trained to identify STO and TSO surface waters using the CARS dataset. Data was extracted from two regional zones within the CARS dataset, selected to represent STO and TSO water (Figure 3). These regions were selected to be representative of the two primary water masses that are known to influence the study region: (A) Coral Sea (STO/EAC) and; (B) Tasman Sea (TSO/non-EAC) waters. Variables extracted from the CARS data set included temperature, salinity and “day of year” (DoY) to account for seasonal variability in water mass characteristics. A total of 58,765 data points were used to train the classification algorithm to identify the two water masses, based on time-varying patterns in their surface temperature and salinity profiles.

(2) The training dataset was then used to fit a logistic regression model to classify water as either STO or TSO. To test how well the model classified unseen portions of the CARS data, the performance of the model was tested using k-fold cross-validation ($k = 10$). This was achieved by randomly partitioning the training dataset into ten equal sized subsamples, using nine of these subsamples to train the model, and holding the remaining one out as a validation set to determine the



accuracy of predictions. This process was then repeated ten times such that each of the subsamples was used once as the validation set. The results of this validation were then averaged to produce an estimation of the model's performance on the CARS data set.

(3) The logistic regression model was then applied to the ROMS model output. The probability of a given cell of water being STO was then calculated by applying the inverse logit function to the log-odds model predictions.


(4) A metric called the "classification ratio" was then calculated from the probability values to explore temporal variability in regional EAC influence, and was defined as the proportion of grid cells in a region classified as STO (probability ≥ 0.5) vs. TSO (probability < 0.5) for each day.

All code was developed with the general purpose programming language "Python" (Rossum, 1995) using the open-source machine learning software library "scikit-learn" (Pedregosa et al., 2011).

3. RESULTS

The classification algorithm produced a metric of EAC presence for the study region and was able to track structural features such as EAC-derived eddies that were evident in the surface temperature and salinity profiles (Figure 4 and Video 1). Model evaluation using k-folds cross-validation on the CARS training data set indicated an accuracy of 99.91%. The method revealed latitudinal variability in the dynamics of the EAC extension, with differences being detected between the three study sites.

The monthly means of the "classification ratio" (ratio of STO to TSO water) demonstrate a sinusoidal seasonal cycle of EAC presence for the three study sites (Figure 5). Furthermore, the annual means of the "classification ratio" demonstrate a positive trend in the ratio of STO water for all three study sites over the 22-year model run. Each study site is characterized by different patterns of EAC dynamics, with mean period of STO dominance




(period of time where the classification ratio is > 0.5) being 246, 207, and 184 days for Jervis Bay, Batemans Bay, and Cape Howe, respectively (**Figure 6**). These patterns also appear to have changed over the study period. When comparing the first and last five years of the dataset, it is apparent that the period of EAC dominance is beginning earlier, persisting longer and ending later for all three study sites (**Figure 6**).

For Jervis Bay the period of EAC dominance has increased by 75 days from 189 to 264 days and is now beginning almost a month earlier and persisting nearly a month later into June. At Batemans Bay the period of EAC dominance has increased by 50 days from 178 to 228 and is now beginning over a month earlier in late October and also persisting for half a month longer. At Cape Howe the period of EAC dominance has increased by 40 days from 156 to 196 days and now begins approximately one month earlier in mid November and ends half a month later at the end of May.

4. DISCUSSION

The goal of this study was to produce a metric of East Australian Current (EAC) presence to assess seasonal and inter-annual patterns of EAC penetration within the EAC extension. This was achieved by developing a supervised machine learning logistic regression algorithm to classify water as either: (A) Sub-tropical origin (STO), representing EAC source water; or (B) Tasman Sea origin (TSO), representing non-EAC water; based on time-varying surface temperature and salinity profiles. The method was able to differentiate spatially complex features of the EAC such as eddies and filaments (**Figure 4**) and identified previously reported patterns of EAC dynamics within the EAC extension, such as a sinusoidal seasonal cycle (Cresswell, 2000), and a trend of intensification over the past 22 years (Ridgway, 2007). In addition, major spatially-explicit changes in the seasonal cycle of EAC dominance were identified at all three study




sites across a 3.5 degree latitudinal gradient of the southeast Australian coastline.

Temporal dynamics of the EAC that were identified using the classification algorithm approach are consistent with that of previous research. The algorithm detected a sinusoidal seasonal cycle in EAC presence for each of the three study sites, with the EAC having a stronger presence in the summer months and a weaker influence during the winter (Ridgway and Godfrey, 1997; Cresswell, 2000). The seasonal cycle of EAC dynamics was found to vary with latitude, with each study site exhibiting different periods of EAC dominance (period of time where the EAC classification ratio was > 0.5). The study sites at more northerly latitudes had a greater mean period of EAC dominance (Figure 6). This was expected as the more northerly sites are closer to the EAC’s separation latitude (between 31 and 33°S) where the majority of the EAC veers eastward as the Tasman Front and the extension zone begins (Cetina-Heredia et al., 2014). Consequently, more northerly sites will be the first to be exposed to the EAC as it penetrates southward during summer, and the last to be exposed as it retreats northward in the winter (Suthers et al., 2011).

Our algorithm also detected a positive inter-annual trend in EAC influence at each of the three study sites between 1994

and 2016 (Figure 5). This is consistent with previous evidence demonstrating a marked increase in the strength, duration, and frequency of southward incursions of EAC water over the last several decades (Holbrook and Bindoff, 1997; Ridgway, 2007; Roemmich et al., 2007; Ridgway et al., 2008). Substantial long-term changes were also detected in the seasonal cycle of EAC presence (Figure 6), with the period of the year being EAC-dominated increasing by approximately 2 months at each of the study sites over the 22-year study period. The onset of the EAC’s dominance now occurs approximately one month earlier and persists for an additional month. The largest change was at the middle site of Batemans Marine Park where the period of the year that is EAC dominated increasing by 75 days from 189 to 264 by the end of the study period (Figure 6).

The encroachment of the EAC’s period of dominance into the winter by 2 months has not previously been reported for the EAC extension and may have significant impacts on marine coastal systems (Figure 6). These changes in EAC dynamics are likely to have flow-on effects to ecological communities through changes in biogeochemical cycling (Oke and Middleton, 2001), alteration of transport of planktonic species and stages (Brierley and Kingsford, 2009; Larsson et al., 2018) and the facilitation of poleward range extension by vagrant tropical species (Ling et al.,



2009). The flow of the EAC influences biogeochemical cycling through upwelling driven by interactions with continental shelf topography (Oke and Middleton, 2001) and eddy generation (Everett et al., 2014). These processes can profoundly influence phytoplankton dynamics by altering the availability of light and nutrients (Ryther, 1969). The shift in the EAC seasonal cycle that we observed in this study could potentially cause an earlier timing of peak spring production of some phytoplankton species, with flow-on effects for the phenology of entire ecological communities (Hallegraeff, 2010; Everett et al., 2014).

The reduced period of marine winter conditions may accelerate the poleward range extension of vagrant tropical species into more temperate regions by allowing them to persist through winter conditions that they would not otherwise be able to survive (Ling et al., 2009; Johnson et al., 2011). This may have implications for the facilitation of community phase shifts from temperate to tropical assemblages (Vergés et al., 2014). Warm-water associated phytoplankton species have undergone range expansions into southeastern Australia, including *Gambierdiscus toxicus* which is now established in southern seagrass beds and can produce harmful algal blooms (Hallegraeff, 2010), and the toxic red-tide dinoflagellate *Noctiluca scintillans* which has expanded its range from Sydney into Southern Tasmanian waters (Hallegraeff, 2010). The increased period of EAC summer dominance may also increase the window of growth of harmful algal bloom taxa. Several well studied dinoflagellates that produce paralytic shellfish toxin, such as

Alexandrium catenella and *Gymnodinium catenatum* bloom in well-defined seasonal temperature windows ($>13^{\circ}\text{C}$ and $>10^{\circ}\text{C}$, respectively) (Hallegraeff et al., 1995; Moore et al., 2009) and will likely see their growth window expand with increasing dominance of the EAC.

Changes in circulation are likely to have effects on dispersal and connectivity patterns for species with pelagic larval stages, and changes in temperature are likely to influence larval survival (Cowen and Sponaugle, 2009; Lett et al., 2010; Byrne et al., 2011). While the alteration of EAC dynamics is likely to have impacts on larval dispersal and survival (Doney et al., 2012) there is evidence that strengthened EAC flow can override the effects of warming on larvae. Cetina-Heredia et al. (2015) showed that while ocean warming can increase the survival of lobster larvae, this was offset by the strengthened EAC current diminishing the supply of larvae to the coast.

These changes to coastal ecosystems may have flow-on effects for the timing of migration and breeding of higher trophic level predators (Johnson et al., 2011). Carroll et al. (2016) observed a reduction in little penguin prey capture success which was coincident with an unusually strong penetration of warm EAC water in the region, mirroring published relationships between commercial sardine catch and SST (John Stewart and Ferrell, 2010; Doubell et al., 2015). The alteration of EAC dynamics may therefore have significant impacts on the foraging capabilities and breeding success of top predator species such as pelagic game fish, marine mammals, and seabirds in this region. As changes

in these predator species' phenology, distribution and trophic and interspecific interactions have the potential to disrupt the ecological function of coastal ecosystems, further work should focus on understanding the influences of changing oceanography on pelagic predators that use coastal waters in this region.

The classification approach outlined in this study is extremely adaptable and may be useful for investigating changes in the dynamics of other variable western boundary current systems. Because the approach only considers the surface properties of the defined water masses alternative data sources such as remotely sensed SST from satellites could be used in place of data from hydrodynamic models. This would extend the suitability of this method to other current systems that may not be covered by any regional models.

Considering the complexity of western boundary current systems, caution should be maintained when interpreting the absolute magnitude of the current classification ratio, and investigations should focus on the changes in current dynamics in the context of previous research and the nuances of the system. Users should carefully consider whether a classification approach is required for their study. For systems that form linear density gradients or can be defined by discrete temperature or salinity values, using metrics such as SST anomaly may be sufficient on their own to investigate watermass dynamics. The classification approach outlined in this study may be advantageous in cases like western boundary currents where watermass properties are highly variable, as it accounts for seasonal time-dependence and uses multiple oceanic variables in its assessment of water-mass presence. By constructing an index of watermass presence probability it is possible to explore the dynamics of western boundary currents in more detail than just investigating the variability in individual ocean properties. To improve the reliability of the approach, future work could explore the use of other physical and biological properties of EAC waters to produce more accurate training and validation datasets. For example, divinyl chlorophyll *a* is a biomarker for tropical Prochlorophytes that is found only in the EAC in this region (Hassler et al., 2011). Although biological indices are more difficult to obtain than physical ocean parameters such as temperature and salinity, they could provide a useful data set to validate the biological relevance of model classifications.

We have developed a classification approach to track highly variable and dynamic ocean features that do not conform to linear mixing models and do not adhere to a linear density gradient or set range of temperature and salinity values. Applying

this method to investigate the temporal dynamics of the EAC extension has revealed significant alterations to the seasonal influence of the EAC for three coastal ecosystems along the southeast coast of Australia. The period of the year that is dominated by the EAC increased by approximately 2 months at each of the study sites between 1994 and 2016. This is a rapid and substantial change in this region, with the potential to significantly alter coastal marine ecosystems within the EAC extension.

DATA AVAILABILITY STATEMENT

The EAC ROMS model output is available at <https://doi.org/10.26190/5e683944e1369> (Kerry and Roughan, 2020a,b). CARS dataset is available at <http://www.marine.csiro.au/~dunn/cars2009/> (Ridgway et al., 2015). All code developed for this study is open-source and hosted online at GitHub as the "Ocean Current Classification" repository (oceancc) (<https://github.com/Lachlan00/oceancc>).

AUTHOR CONTRIBUTIONS

All authors contributed to this paper. Study design: LP, IJ, RH, GC, and MR. Data collection and processing: LP and MR. Analysis: LP and IJ. The paper was written by LP but benefited from input from all authors.

FUNDING

This study was funded by Australian Research Council Linkage Grants (LP110200603 and LP160100162) with contributions from the Taronga Conservation Society Australia.

ACKNOWLEDGMENTS

We thank the two reviewers and Dr. Colette Kerry whose comments greatly improved the quality of the manuscript. We would also like to thank Dr. Colette Kerry for her assistance in obtaining and processing the data from the ROMS hydrodynamic model.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmars.2020.00365/full#supplementary-material>

REFERENCES

- Archer, M. R., Keating, S. R., Roughan, M., Johns, W. E., Lumpkin, R., Beron-Vera, F. J., et al. (2018). The kinematic similarity of two western boundary currents revealed by sustained high-resolution observations. *Geophys. Res. Lett.* 45, 6176–6185. doi: 10.1029/2018GL078429
- Archer, M. R., Roughan, M., Keating, S. R., and Schaeffer, A. (2017). On the variability of the East Australian Current: jet structure, meandering, and influence on shelf circulation. *J. Geophys. Res.* 122, 8464–8481. doi: 10.1002/2017JC013097
- Baird, M. E., Suthers, I. M., Griffin, D. A., Hollings, B., Pattiaratchi, C., Everett, J. D., et al. (2011). The effect of surface flooding on the physical-biogeochemical dynamics of a warm-core eddy off southeast Australia. *Deep Sea Res. Part II* 58, 592–605. doi: 10.1016/j.dsr2.2010.10.002
- Brassington, G. B., Summons, N., and Lumpkin, R. (2011). Observed and simulated lagrangian and eddy characteristics of the East Australian

- Current and the Tasman Sea. *Deep Sea Res. Part II* 58, 559–573. doi: 10.1016/j.dsr2.2010.10.001
- Brierley, A. S., and Kingsford, M. J. (2009). Impacts of climate change on marine organisms and ecosystems. *Curr. Biol.* 19, R602–R614. doi: 10.1016/j.cub.2009.05.046
- Bull, C. Y. S., Kiss, A. E., Jourdain, N. C., England, M. H., and van Sebille, E. (2017). Wind forced variability in eddy formation, eddy shedding, and the separation of the East Australian Current. *J. Geophys. Res.* 122, 9980–9998. doi: 10.1002/2017JC013311
- Byrne, M., Selvakumaraswamy, P., Ho, M., Woolsey, E., and Nguyen, H. (2011). Sea urchin development in a global change hotspot, potential for southerly migration of thermotolerant propagules. *Deep Sea Res. Part II* 58, 712–719. doi: 10.1016/j.dsr2.2010.06.010
- Cai, W. (2006). Antarctic ozone depletion causes an intensification of the southern ocean super-gyre circulation. *Geophys. Res. Lett.* 33, 720–738. doi: 10.1029/2005GL024911
- Cai, W., Shi, G., Cowan, T., Bi, D., and Ribbe, J. (2005). The response of the southern annular mode, the East Australian Current, and the southern mid-latitude ocean circulation to global warming. *Geophys. Res. Lett.* 32, 136–148. doi: 10.1029/2005GL024701
- Carroll, G., Cox, M., Harcourt, R., Pitcher, B. J., Slip, D., and Jonsen, I. (2017). Hierarchical influences of prey distribution on patterns of prey capture by a marine predator. *Funct. Ecol.* 31, 1750–1760. doi: 10.1111/1365-2435.12873
- Carroll, G., Everett, J. D., Harcourt, R., Slip, D., and Jonsen, I. (2016). High sea surface temperatures driven by a strengthening current reduce foraging success by penguins. *Sci. Rep.* 6:22236. doi: 10.1038/srep22236
- Cetina-Heredia, P., Roughan, M., Sebille, E., Feng, M., and Coleman, M. A. (2015). Strengthened currents override the effect of warming on lobster larval dispersal and survival. *Glob. Change Biol.* 21, 4377–4386. doi: 10.1111/gcb.13063
- Cetina-Heredia, P., Roughan, M., van Sebille, E., and Coleman, M. A. (2014). Long-term trends in the East Australian Current separation latitude and eddy driven transport. *J. Geophys. Res.* 119, 4351–4366. doi: 10.1002/2014JC010071
- Cowen, R. K., and Spoungue, S. (2009). Larval dispersal and marine population connectivity. *Annu. Rev. Mar. Sci.* 1, 443–466. doi: 10.1146/annurev.marine.010908.163757
- Cresswell, G. R. (1982). The coalescence of two East Australian Current warm-core eddies. *Science* 215, 161–164. doi: 10.1126/science.215.4529.161
- Cresswell, G. R. (1983). Physical evolution of Tasman Sea eddy. *J. Mar. Freshw. Res.* 34, 495–513. doi: 10.1071/MF9830495
- Cresswell, G. R. (2000). Currents of the continental shelf and upper slope of Tasmania. *Pap. Proc. R. Soc. Tasmania* 133, 21–30. doi: 10.26749/rstpp.133.3.21
- Doney, S. C., Ruckelshaus, M., Duffy, J. E., Barry, J. P., Chan, F., English, C. A., et al. (2012). Climate change impacts on marine ecosystems. *Annu. Rev. Mar. Sci.* 4, 11–37. doi: 10.1146/annurev-marine-041911-111611
- Doubell, M., Ward, T., Watson, P., James, C., Carroll, J., and Redondo Rodriguez, A. (2015). *Optimising the Size and Quality of Sardines Through Real-Time Harvesting*. Technical Report 2013/746, South Australian Research and Development Institute (Aquatic Sciences), Adelaide.
- Eddyane, K. S. (2003). *Conservation, Monitoring and Recovery of Threatened Giant Kelp (Macrocystis pyrifera) Forests in Tasmania*. Report to Environment Australia (Marine Species Protection Program). Department of Primary Industries, Water and Environment, Hobart.
- Everett, J. D., Baird, M. E., Roughan, M., Suthers, I. M., and Doblin, M. A. (2014). Relative impact of seasonal and oceanographic drivers on surface chlorophyll a along a western boundary current. *Prog. Oceanogr.* 120, 340–351. doi: 10.1016/j.pocean.2013.10.016
- Gangopadhyay, A., Rezendes, J. A., Lydon, K., Balasubramanian, R., and Valova, I. (2015). “Towards an automated detection of the Gulf stream north wall from concurrent satellite images,” in *OCEANS 2015 - MTS/IEEE Washington* (Washington, DC), 1–8. doi: 10.23919/OCEANS.2015.7404566
- Godfrey, J. S., Cresswell, G. R., Golding, T. J., Pearce, A. F., and Boyd, R. (1980). The separation of the East Australian Current. *J. Phys. Oceanogr.* 10, 430–440. doi: 10.1175/1520-0485(1980)010<0430:TSOTEA>2.0.CO;2
- Hallegraef, G., McCausland, M., and Brown, R. (1995). Early warning of toxic dinoflagellate blooms of gymnodinium catenatum in southern Tasmanian waters. *J. Plankt. Res.* 17, 1163–1176. doi: 10.1093/plankt/17.6.1163
- Hallegraef, G. M. (2010). Ocean climate change, phytoplankton community responses, and harmful algal blooms: a formidable predictive challenge. *J. Phycol.* 46, 220–235. doi: 10.1111/j.1529-8817.2010.00815.x
- Harris, G., Griffiths, F., Clementson, L., Lyne, V., and Doe, H. D. (1991). Seasonal and interannual variability in physical processes, nutrient cycling and the structure of the food chain in Tasmanian shelf waters. *J. Plankton Res.* 13(Suppl. 1), 109–131.
- Harris, G., Nilsson, C., Clementson, L., and Thomas, D. (1987). The water masses of the east coast of Tasmania: seasonal and interannual variability and the influence on phytoplankton biomass and productivity. *Mar. Freshw. Res.* 38, 569–590. doi: 10.1071/MF9870569
- Hassler, C., Djajadikarta, J., Doblin, M., Everett, J., and Thompson, P. (2011). Characterisation of water masses and phytoplankton nutrient limitation in the East Australian Current separation zone during spring 2008. *Deep Sea Res. Part II* 58, 664–677. doi: 10.1016/j.dsr2.2010.06.008
- Hobday, A., Young, J., Moeseneder, C., and Dambacher, J. (2011). Defining dynamic pelagic habitats in oceanic waters off eastern Australia. *Deep Sea Res. Part II* 58, 734–745. doi: 10.1016/j.dsr2.2010.10.006
- Hogg, N. G., and Johns, W. E. (1995). Western boundary currents. *Rev. Geophys.* 33, 1311–1334. doi: 10.1029/95RG00491
- Holbrook, N. J., and Bindoff, N. L. (1997). Interannual and decadal temperature variability in the southwest Pacific Ocean between 1955 and 1988. *J. Clim.* 10, 1035–1049. doi: 10.1175/1520-0442(1997)010<1035:IADTVI>2.0.CO;2
- Hosmer, D. W. Jr., Lemeshow, S., and Sturdivant, R. X. (2013). *Applied Logistic Regression, Vol. 398*. Hoboken, NJ: John Wiley & Sons. doi: 10.1002/9781118548387
- Imawaki, S., Bower, A. S., Beal, L., and Qiu, B. (2013). “Chapter 13 - western boundary currents,” in *Ocean Circulation and Climate, Vol. 103 of International Geophysics*, eds G. Siedler, S. M. Griffies, J. Gould, and J. A. Church (Kidlington, UK: Academic Press), 305–338. doi: 10.1016/B978-0-12-391851-2.00013-1
- John Stewart, G. B., and Ferrell, D. (2010). *Review of the Biology and Fishery for Australian Sardines (Sardinops sagax) in New South Wales*. Fisheries research report series, Industry & Investment NSW.
- Johnson, C. R., Banks, S. C., Barrett, N. S., Cazassus, F., Dunstan, P. K., Edgar, G. J., et al. (2011). Climate change cascades: Shifts in oceanography, species’ ranges and subtidal marine community dynamics in eastern Tasmania. *J. Exp. Mar. Biol. Ecol.* 400, 17–32. doi: 10.1016/j.jembe.2011.02.032
- Jones, C. T., Sikora, T. D., Vachon, P. W., and Buckley, J. R. (2014). Ocean feature analysis using automated detection and classification of sea-surface temperature front signatures in radarsat-2 images. *Bull. Am. Meteorol. Soc.* 95, 677–679. doi: 10.1175/BAMS-D-12-00174.1
- Kelly, P., Clementson, L., Davies, C., Corney, S., and Swadling, K. (2016). Zooplankton responses to increasing sea surface temperatures in the southeastern Australia global marine hotspot. *Estuar. Coast. Shelf Sci.* 180, 242–257. doi: 10.1016/j.ecss.2016.07.019
- Kerry, C., and Roughan, M. (2020a). Downstream evolution of the East Australian Current system: mean flow, seasonal and intra-annual variability. *J. Geophys. Res.* e2019JC015227. doi: 10.1029/2019JC015227
- Kerry, C., and Roughan, M. (2020b). A high-resolution, 22-year, free-running, hydrodynamic simulation of the East Australian Current system using the regional ocean modeling system. *UNSW.dataset*. doi: 10.26190/5e683944e1369
- Kotsiantis, S. B. (2007). “Supervised machine learning: a review of classification techniques,” in *Proceedings of the 2007 Conference on Emerging Artificial Intelligence Applications in Computer Engineering: Real World AI Systems with Applications in eHealth HCI, Information Retrieval and Pervasive Technologies* (Amsterdam: IOS Press), 3–24.
- Larsson, M., Laczka, O., Suthers, I., Ajani, P., and Doblin, M. (2018). Hitchhiking in the East Australian Current: rafting as a dispersal mechanism for harmful epibenthic dinoflagellates. *Mar. Ecol. Prog. Ser.* 596, 49–60. doi: 10.3354/meps12579
- Lett, C., Ayata, S.-D., Huret, M., and Irisson, J.-O. (2010). Biophysical modelling to investigate the effects of climate change on marine population dispersal and connectivity. *Prog. Oceanogr.* 87, 106–113. doi: 10.1016/j.pocean.2010.09.005
- Ling, S. D., Johnson, C. R., Ridgway, K., Hobday, A. J., and Haddon, M. (2009). Climate-driven range extension of a sea urchin: inferring future trends by analysis of recent population dynamics. *Glob. Change Biol.* 15, 719–731. doi: 10.1111/j.1365-2486.2008.01734.x

- Moore, S. K., Mantua, N. J., Hickey, B. M., and Trainer, V. L. (2009). Recent trends in paralytic shellfish toxins in puget sound, relationships to climate, and capacity for prediction of toxic events. *Harmful Algae* 8, 463–477. doi: 10.1016/j.hal.2008.10.003
- Mos, B., Ah Yong, S. T., Burnes, C. N., Davie, P. J. F., and McCormack, R. B. (2017). Range extension of a euryhaline crab, *Varuna litterata* (fabricius, 1798) (brachyura: Varunidae), in a climate change hot-spot. *J. Crustacean Biol.* 37, 258–262. doi: 10.1093/jcbiol/rux030
- Nilsson, C., and Cresswell, G. (1980). The formation and evolution of East Australian Current warm-core eddies. *Prog. Oceanogr.* 9, 133–183. doi: 10.1016/0079-6611(80)90008-7
- Oke, P. R., Brassington, G. B., Griffin, D. A., and Schiller, A. (2008). The blueslink ocean data assimilation system (bodas). *Ocean Model.* 21, 46–70. doi: 10.1016/j.ocemod.2007.11.002
- Oke, P. R., and Middleton, J. H. (2001). Nutrient enrichment off Port Stephens: the role of the East Australian Current. *Contin. Shelf Res.* 21, 587–606. doi: 10.1016/S0278-4343(00)00127-8
- Oliver, E. C. J., Benthuisen, J. A., Bindoff, N. L., Hobday, A. J., Holbrook, N. J., Mundy, C. N., et al. (2017). The unprecedented 2015/16 Tasman Sea marine heatwave. *Nat. Commun.* 8:16101. doi: 10.1038/ncomms16101
- Oliver, E. C. J., and Holbrook, N. J. (2014). Extending our understanding of South Pacific gyre “spin-up”: Modeling the East Australian Current in a future climate. *J. Geophys. Res.* 119, 2788–2805. doi: 10.1002/2013JC009591
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., et al. (2011). Scikit-learn: Machine learning in python. *J. Mach. Learn. Res.* 12, 2825–2830. doi: 10.5555/1953048.2078195
- Phillips, L. R., Hindell, M., Hobday, A. J., and Lea, M.-A. (2019). Variability in at-sea foraging behaviour of little penguins *eudyptula minor* in response to finescale environmental features. *Mar. Ecol. Prog. Ser.* 627, 141–154. doi: 10.3354/meps13095
- Pittock, B. (2003). *Climate Change: An Australian Guide to the Science and Potential Impacts*. Canberra ACT: Australian Greenhouse Office.
- Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., and Schlax, M. G. (2007). Daily high-resolution-blended analyses for sea surface temperature. *J. Clim.* 20, 5473–5496. doi: 10.1175/2007JCLI1824.1
- Ridgway, K., and Dunn, J. (2003). Mesoscale structure of the mean East Australian Current system and its relationship with topography. *Prog. Oceanogr.* 56, 189–222. doi: 10.1016/S0079-6611(03)00004-1
- Ridgway, K., Dunn, J., and Wilkin, J. (2015). *Cars 2009 - CSIRO Atlas of Regional Seas - World Monthly*. Argo - Broad scale Global Array of Temperature/Salinity Profiling Floats.
- Ridgway, K. R. (2007). Long-term trend and decadal variability of the southward penetration of the East Australian Current. *Geophys. Res. Lett.* 34, 345–365. doi: 10.1029/2007GL030393
- Ridgway, K. R., Coleman, R. C., Bailey, R. J., and Sutton, P. (2008). Decadal variability of East Australian Current transport inferred from repeated high-density XBT transects, a CTD survey and satellite altimetry. *J. Geophys. Res.* 113, 346–365. doi: 10.1029/2007JC004664
- Ridgway, K. R., Dunn, J. R., and Wilkin, J. L. (2002). Ocean interpolation by four-dimensional weighted least squares - application to the waters around Australasia. *J. Atmos. Ocean. Technol.* 19:1357. doi: 10.1175/1520-0426(2002)019<1357:OIBFDW>2.0.CO;2
- Ridgway, K. R., and Godfrey, J. S. (1997). Seasonal cycle of the East Australian Current. *J. Geophys. Res.* 102, 22921–22936. doi: 10.1029/97JC00227
- Roemmich, D., Gilson, J., Davis, R., Sutton, P., Wijffels, S., and Riser, S. (2007). Decadal spinup of the South Pacific subtropical gyre. *J. Phys. Oceanogr.* 37, 162–173. doi: 10.1175/JPO3004.1
- Rossum, G. (1995). *Python Reference Manual*. Technical Report, Python Software Foundation, Amsterdam.
- Roughan, M., Macdonald, H. S., Baird, M. E., and Glasby, T. M. (2011). Modelling coastal connectivity in a western boundary current: Seasonal and inter-annual variability. *Deep Sea Res. Part II* 58, 628–644. doi: 10.1016/j.dsr2.2010.06.004
- Ryther, J. H. (1969). Photosynthesis and fish production in the sea. *Science* 166, 72–76. doi: 10.1126/science.166.3901.72
- Shchepetkin, A. F., and McWilliams, J. C. (2003). A method for computing horizontal pressure-gradient force in an oceanic model with a nonaligned vertical coordinate. *J. Geophys. Res.* 108, FET 35-1–FET 35-34. doi: 10.1029/2001JC001047
- Shchepetkin, A. F., and McWilliams, J. C. (2005). The regional oceanic modeling system (roms): a split-explicit, free-surface, topography-following-coordinate oceanic model. *Ocean Model.* 9, 347–404. doi: 10.1016/j.ocemod.2004.08.002
- Suthers, I. M., Young, J. W., Baird, M. E., Roughan, M., Everett, J. D., Brassington, G. B., et al. (2011). The strengthening East Australian Current, its eddies and biological effects – an introduction and overview. *Deep Sea Res. Part II* 58, 538–546. doi: 10.1016/j.dsr2.2010.09.029
- Thompson, P., Baird, M., Ingleton, T., and Doblin, M. (2009). Long-term changes in temperate Australian coastal waters: implications for phytoplankton. *Mar. Ecol. Prog. Ser.* 394, 1–19. doi: 10.3354/meps08297
- Thresher, R., Proctor, C., Ruiz, G., Gurney, R., MacKinnon, C., Walton, W., et al. (2003). Invasion dynamics of the European shore crab, *Carcinus maenas*, in Australia. *Mar. Biol.* 142, 867–876. doi: 10.1007/s00227-003-1011-1
- Tilburg, C. E., Hurlburt, H. E., O'Brien, J. J., and Shriver, J. F. (2001). The dynamics of the East Australian Current system: The Tasman front, the east Auckland current, and the east cape current. *J. Phys. Oceanogr.* 31, 2917–2943. doi: 10.1175/1520-0485(2001)031<2917:TDOTEA>2.0.CO;2
- Vergés, A., Steinberg, P. D., Hay, M. E., Poore, A. G. B., Campbell, A. H., Ballesteros, E., et al. (2014). The tropicalization of temperate marine ecosystems: climate-mediated changes in herbivory and community phase shifts. *Proc. R. Soc. B* 281:20140846. doi: 10.1098/rspb.2014.0846
- Weiss, S. M., and Kulikowski, C. A. (1991). *Computer Systems That Learn: Classification and Prediction Methods from Statistics, Neural Nets, Machine Learning, and Expert Systems*. San Francisco, CA: Morgan Kaufmann Publishers Inc.
- Wu, L., Cai, W., Zhang, L., Nakamura, H., Timmermann, A., Joyce, T., et al. (2012). Enhanced warming over the global subtropical western boundary currents. *Nat. Clim. Change* 2, 161–166. doi: 10.1038/nclimate1353
- Yu, L., and Weller, R. A. (2007). Objectively analyzed air-sea heat fluxes for the global ice-free oceans (1981–2005). *Bull. Am. Meteorol. Soc.* 88, 527–540. doi: 10.1175/BAMS-88-4-527

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Phillips, Carroll, Jonsen, Harcourt and Roughan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.